

$N = 2 \rightarrow N = 1$ supergravity reduction in four dimensions

Laura Andrianopoli ¹

¹ Dip. Fisica, Politecnico di Torino, Corso Duca degli Abruzzi 24, I-10129 Torino
e-mail: andrianopoli@athena.polito.it

Abstract: We discuss the reduction of $N = 2$ supergravity to $N = 1$, by a consistent truncation of the second gravitino multiplet.

It is well known that for globally supersymmetric theories with particle content of spin $0, \frac{1}{2}, 1$ any theory with N supersymmetries can be regarded as a particular case of a theory with a number $N' < N$ of supersymmetries [1]. To prove this it is sufficient to decompose the N supersymmetry-extended multiplets into N' -multiplets. Of course N -extended supersymmetry is more restrictive than $N' < N$ supersymmetry implying that the former will only allow some restricted couplings of the latter. As we are going to show, the same argument does not apply to supergravity theories [2]. Indeed, let us consider a standard N -extended supergravity with N gravitini and a given number of matter multiplets: then the N' -extended supergravity obtained by reduction from the mother theory will no longer be standard because a certain number $N - N'$ of spin $\frac{3}{2}$ multiplets appear in the decomposition. To obtain a standard N' -extended supergravity one must truncate out at least the $N - N'$ spin $\frac{3}{2}$ multiplets and all the non-linear couplings that they generate in the supergravity action.

Here I will report on the truncation of $N = 2$ matter-coupled supergravity theory to $N = 1$, showing that the reduction does indeed imply the truncation of part of the matter content, besides of the second gravitino multiplet. The present discussion is based on [2]. A more detailed analysis, with proofs of consistency, and including the truncation of $N = 8$ supergravity down to general N theories can be found in [2] to which I refer also for the notations and conventions and for a complete list of references.

The supersymmetry reduction $N = 2 \rightarrow N = 1$ is obtained by truncating the $N = 1$ spin $3/2$ multiplet containing the second gravitino $\psi_{\mu 2}$ and the graviphoton.

Let us write down the supersymmetry transformation laws of the fermions for the $N = 2$ theory, up to 3-fermions terms [3]:

$$\delta \psi_{A\mu} = \hat{\nabla}_\mu \epsilon_A + \left(i g S_{AB} \eta_{\mu\nu} + \epsilon_{AB} T_{\mu\nu}^- \right) \gamma^\nu \epsilon^B \quad (1)$$

$$\delta \lambda^{\mathcal{I}A} = i \nabla_\mu z^{\mathcal{I}} \gamma^\mu \epsilon^A + G_{\mu\nu}^{-\mathcal{I}} \gamma^{\mu\nu} \epsilon_B \epsilon^{AB} + g W^{\mathcal{I}AB} \epsilon_B \quad (2)$$

$$\delta \zeta_\alpha = i \mathcal{U}_u^{B\beta} \nabla_\mu q^u \gamma^\mu \epsilon^A \epsilon_{AB} C_{\alpha\beta} + g N_\alpha^A \epsilon_A \quad (3)$$

where: $\hat{\nabla}_\mu \epsilon_A = \mathcal{D}_\mu \epsilon_A + \hat{\omega}_{\mu|A}{}^B \epsilon_B + \hat{\mathcal{Q}}_\mu \epsilon_A$, with the $SU(2)$ and $U(1)$ 1-form “gauged” connections respectively given by: $\hat{\omega}_A{}^B = \omega_A{}^B + g_{(\Lambda)} A^\Lambda P_\Lambda^x (\sigma^x)_A{}^B$; $\hat{\mathcal{Q}} = \mathcal{Q} + g_{(\Lambda)} A^\Lambda P_\Lambda^0$,

$\omega_A{}^B$ and $\mathcal{Q} = -\frac{i}{2}(\partial_{\mathcal{I}}\mathcal{K}dz^{\mathcal{I}} - \partial_{\bar{\mathcal{I}}}\mathcal{K}d\bar{z}^{\bar{\mathcal{I}}})$ being the $SU(2)$ and $U(1)$ connections of the ungauged theory. $T_{\mu\nu}^-$ appearing in the supersymmetry transformation law of the $N = 2$ left-handed gravitini is the “dressed” graviphoton defined as:

$$T_{\mu\nu}^- \equiv 2i\text{Im}\mathcal{N}_{\Lambda\Sigma}L^{\Sigma}\left[F_{\mu\nu}^{\Lambda-} + 3 \text{ fermions terms}\right] \quad (4)$$

while

$$G_{\mu\nu}^{\mathcal{I}-} \equiv -g^{\mathcal{I}\bar{\mathcal{J}}}\text{Im}\mathcal{N}_{\Lambda\Sigma}f_{\bar{\mathcal{J}}}^{\Sigma}\left[F_{\mu\nu}^{\Lambda-} + 3 \text{ fermions terms}\right] \quad (5)$$

are the “dressed” field strengths of the vectors inside the vector multiplets (the “minus” apex means taking the self-dual part.). Moreover the fermionic shifts S_{AB} , $W^{\mathcal{I}AB}$ and N_{α}^A are given in terms of the prepotentials and Killing vectors of the quaternionic geometry as follows:

$$S_{AB} = i\frac{1}{2}P_{AB\Lambda}L^{\Lambda} \equiv i\frac{1}{2}P_{\Lambda}^x\sigma_{AB}^xL^{\Lambda} \quad (6)$$

$$W^{\mathcal{I}AB} = iP_{\Lambda}^{AB}g^{\mathcal{I}\bar{\mathcal{J}}}f_{\bar{\mathcal{J}}}^{\Lambda} + \epsilon^{AB}k_{\Lambda}^{\mathcal{I}}\bar{L}^{\Lambda} \quad (7)$$

$$N_{\alpha}^A = 2\mathcal{U}_{\alpha u}^A k_{\Lambda}^u \bar{L}^{\Lambda} \quad (8)$$

Since we are going to compare the $N = 2$ reduced theory with the standard $N = 1$ supergravity, I also quote the supersymmetry transformation laws of fermions in the latter theory [4],[5]. We have, up to 3-fermions terms:

$N = 1$ transformation laws

$$\delta\psi_{\bullet\mu} = \mathcal{D}_{\mu}\epsilon_{\bullet} + \frac{i}{2}\hat{Q}_{\mu}\epsilon_{\bullet} + iL(z, \bar{z})\gamma_{\mu}\varepsilon^{\bullet} \quad (9)$$

$$\delta\chi^i = i\left(\partial_{\mu}z^i + g_{(\Lambda)}A_{\mu}^{\Lambda}k_{\Lambda}^i\right)\gamma^{\mu}\varepsilon_{\bullet} + N^i\varepsilon_{\bullet} \quad (10)$$

$$\delta\lambda_{\bullet}^{\Lambda} = \mathcal{F}_{\mu\nu}^{(-)\Lambda}\gamma^{\mu\nu}\varepsilon_{\bullet} + iD^{\Lambda}\varepsilon_{\bullet} \quad (11)$$

where $\hat{Q}$ is defined in a way analogous to the $N = 2$ definition and:

$$L(z, \bar{z}) = W(z)e^{\frac{1}{2}\mathcal{K}_{(1)}(z, \bar{z})}, \quad \nabla_{\bar{i}}L = 0 \quad (12)$$

$$N^i = 2g^{i\bar{j}}\nabla_{\bar{j}}\bar{L} \quad (13)$$

$$D^{\Lambda} = -2(\text{Im}f_{\Lambda\Sigma})^{-1}P_{\Sigma}(z, \bar{z}) \quad (14)$$

and $W(z), \mathcal{K}_{(1)}(z, \bar{z}), P_{\Sigma}(z, \bar{z}), f_{\Lambda\Sigma}(z)$ are the superpotential, Kähler potential, Killing prepotential and vector kinetic matrix respectively [5], [4], [6]. Note that for the gravitino and gaugino fields we have denoted by a lower (upper) dot left-handed (right-handed) chirality. For the spinors of the chiral multiplets χ , instead, left-handed (right-handed) chirality is encoded via an upper holomorphic (antiholomorphic) world index ($\chi^i, \bar{\chi}^{\bar{i}}$).

To perform the truncation we set $A=1$ and 2 successively, putting $\psi_{2\mu} = \epsilon_2 = 0$, and for the gravitino we get, from equation (1):

$$\delta\psi_{1\mu} = \mathcal{D}_{\mu}\epsilon_1 - \hat{Q}_{\mu}\epsilon_1 - \hat{\omega}_{\mu|1}{}^1\epsilon_1 + igS_{11}\eta_{\mu\nu}\gamma^{\nu}\epsilon^1 \quad (15)$$

while, for consistency:

$$\delta\psi_{2\mu} \equiv 0 = -\hat{\omega}_{\mu|2}{}^1\epsilon_1 + (igS_{21}\eta_{\mu\nu} - T_{\mu\nu}^-)\gamma^{\nu}\epsilon^1 \quad (16)$$

Comparing (9) with (15), we learn that we must identify $\psi_{1\mu} \equiv \psi_{\bullet\mu}$; $\epsilon_1 \equiv \epsilon_{\bullet}$. Furthermore, gS_{11} must be identified with the superpotential of the $N = 1$ theory, that is to

the covariantly holomorphic section L of the $N = 1$ Kähler-Hodge manifold. Therefore we have [7] $L(q, z, \bar{z}) = \frac{i}{2} g_{(\Lambda)} P_{\Lambda}^x (\sigma^x)_{11} L^{\Lambda} = \frac{i}{2} g_{(\Lambda)} (P_{\Lambda}^1 - i P_{\Lambda}^2) L^{\Lambda}$. As it has been shown in [2], after consistent reduction of the special-Kähler manifold $\mathcal{M}^{SK}$ and of the quaternionic σ -model $\mathcal{M}^Q$, L is in fact a covariantly holomorphic function of the Kähler coordinates w^s of the reduced manifold $\mathcal{M}^{KH} \subset \mathcal{M}^Q$ and of some subset $z^i \in \mathcal{M}_R$ of the scalars $z^{\mathcal{I}}$ of the $N = 2$ special-Kähler manifold $\mathcal{M}^{SK}$.

Furthermore, for a consistent truncation we must set to zero all the following bosonic structures:

$$T_{\mu\nu}^- = 0 \quad (17)$$

$$\hat{\omega}_{\mu|2}{}^1 = 0 \quad (18)$$

$$S_{21} = 0 \quad (19)$$

As it has been proven in [2], equations (17) and (18) are “orthogonality” conditions on the scalar sectors of $N = 2$ supergravity which imply a reduction of both special Kähler manifold ($\mathcal{M}^{SK}(n_V)$) [8],[9], [10], [3] and quaternionic manifold ($\mathcal{M}^Q(n_H)$), where n_V and n_H are the number of vector multiplets and hypermultiplets respectively, and imply therefore a truncation of part of the matter multiplets. We note that similar orthogonality conditions, leading to the same reduction of the matter content of the theory, can be found through a complementary analysis, by looking at the 3-fermions terms in the transformation laws, instead of at the bosonic ones [11].

Equation (17) is satisfied by imposing a suitable constraint on the set of vectors and of scalar sections which can be retained in the reduction. Indeed, if we decompose the index Λ labelling the vectors into two disjoint sets $\Lambda \Rightarrow (\Lambda, X)$, $\Lambda = 1, \dots, n'_V = n_V - n_C$; $X = 0, 1, \dots, n_C$, we may satisfy the relation (17) as an “orthogonality relation” between the subset Λ running on the retained vectors and the subset X running on the retained scalar sections. That is we set:

$$F_{\mu\nu}^X = 0; \quad (20)$$

$$\text{Im} \mathcal{N}_{\Lambda\Sigma} L^{\Sigma} = T_{\Lambda} = 0 \quad (21)$$

We note that if we delete the electric field strengths F^{-X} we must also delete their magnetic counterpart $\mathcal{G}_{\bar{X}} = \bar{\mathcal{N}}_{XY} F^{-Y} + \bar{\mathcal{N}}_{X\Sigma} F^{-\Sigma} = 0$ so that we must also impose $\mathcal{N}_{X\Sigma} = 0$. Then, the constraint (21) reduces to $\text{Im} \mathcal{N}_{\Lambda\Sigma} L^{\Sigma} = 0$ which implies

$$L^{\Sigma} = 0 \quad (22)$$

since the vector-kinetic matrix $\text{Im} \mathcal{N}_{\Lambda\Sigma}$ has to be invertible. Note that condition (22) implies a reduction of the $N = 2$ scalar manifold $\mathcal{M}^{SK} \rightarrow \mathcal{M}_R$, since it says that some coordinate dependent sections on $\mathcal{M}^{SK}$ have to be zero in the reduced theory.

Let us now quote the reduction of the gauginos transformation law. When $\epsilon_2 = 0$ we get, up to three fermions terms:

$$\delta \lambda^{\mathcal{I}1} = i \nabla_{\mu} z^{\mathcal{I}} \gamma^{\mu} \epsilon^1 + W^{\mathcal{I}11} \epsilon_1 \quad (23)$$

$$\delta \lambda^{\mathcal{I}2} = -G_{\mu\nu}^{-\mathcal{I}} \gamma^{\mu\nu} \epsilon_1 + g W^{\mathcal{I}21} \epsilon_1. \quad (24)$$

From eqs. (23) and (24) we immediately see that the spinors $\lambda^{\mathcal{I}1}$ transform into the scalars $z^{\mathcal{I}}$ (and should therefore belong to $N = 1$ chiral multiplets) while the spinors $\lambda^{\mathcal{I}2}$ transform into the matter vectors field strengths $G_{\mu\nu}^{-\mathcal{I}}$ (and should then be identified with the gauginos of the $N = 1$ vector multiplets). Let us decompose the world indices

$\mathcal{I}$ of the $N = 2$ special-Kähler σ -model as follows: $\mathcal{I} \Rightarrow (i, \alpha)$, with $i = 1, \dots, n_C$, $\alpha = 1, \dots, n'_V = n_V - n_C$, where n_C and n'_V are respectively the number of chiral and vector multiplets of the reduced $N = 1$ theory while n_V is the number of $N = 2$ vector multiplets. By an appropriate choice of coordinates, we call z^i the coordinates on $\mathcal{M}_R$, z^α the coordinates on the orthogonal complement. Then it is easy to see that the metric $g_{\mathcal{I}\bar{\mathcal{J}}}$ has only components $g_{i\bar{j}}, g_{\alpha\bar{\beta}}$, while $g_{i\bar{\alpha}} = 0$. Then, if we decompose the gauginos $\lambda^{\mathcal{I}2} \Rightarrow (\lambda^{i2}, \lambda^{\alpha 2})$, the above truncation implies, by supersymmetry, $\lambda^{i2} = 0$ and, for consistency,

$$\delta \lambda^{i2} = -G_{\mu\nu}^{-i} \gamma^{\mu\nu} \epsilon_1 + g W^{i21} \epsilon_1 = 0. \quad (25)$$

Setting $G_{\mu\nu}^{-i} = 0$ gives:

$$G_{\mu\nu}^{-i} = -g^{i\bar{j}} \nabla_{\bar{j}} \bar{L}^\Lambda \text{Im} \mathcal{N}_{\Lambda\Sigma} F_{\mu\nu}^{-\Sigma} = -g^{i\bar{j}} \nabla_{\bar{j}} \bar{L}^\Lambda \text{Im} \mathcal{N}_{\Lambda\Sigma} F_{\mu\nu}^{-\Sigma} = 0 \quad (26)$$

implying

$$\nabla_{\bar{j}} \bar{L}^\Lambda = \bar{f}_{\bar{j}}^\Lambda = 0. \quad (27)$$

Moreover, $W^{i21} = 0$ implies $P_\Lambda^3 = 0$, $k_\Lambda^i = 0$. Note that the integrability condition of equation (27) is:

$$\nabla_i \nabla_j L^\Lambda = i C_{ijk} g^{\mathcal{K}\bar{\mathcal{K}}} \nabla_{\bar{\mathcal{K}}} \bar{L}^\Lambda = i C_{ijk} g^{k\bar{k}} \nabla_{\bar{k}} \bar{L}^\Lambda + i C_{ij\alpha} g^{\alpha\bar{\alpha}} \nabla_{\bar{\alpha}} \bar{L}^\Lambda = 0. \quad (28)$$

where C_{ijk} is the 3-index symmetric tensor appearing in the equations defining the special geometry (see *e.g.* ref. [12],[3]). Since the first term on the r.h.s. of equation (28) is zero on $\mathcal{M}_R$ (equation (27)), equation (28) is satisfied by imposing:

$$C_{ij\alpha} = 0 \quad (29)$$

so that only the $N = 2$ special-Kähler manifolds satisfying the constraint (29) are suitable for reduction.

Let us now consider the other condition coming from the reduction of the gravitino transformation law, eq. (18). It implies, on the curvature Ω_2^1 of the $SU(2)$ connection ω_2^1 :

$$\Omega_2^1 = -i \mathbb{C}_{\alpha\beta} \mathcal{U}_1^\alpha \wedge \mathcal{U}^{\beta 2} = 0 \quad (30)$$

Let us now note that the scalars in $N = 1$ supergravity must lie in chiral multiplets, and span in general a Kähler-Hodge manifold. It is therefore required that the holonomy of the quaternionic manifold be reduced: $\text{Hol}(\mathcal{M}^Q) \subset SU(2) \times Sp(2n_H) \rightarrow \text{Hol}(\mathcal{M}^{KH}) \subset U(1) \times SU(n)$. Therefore the $SU(2)$ and the $Sp(2n_H)$ indices have to be decomposed accordingly. We set $\alpha \rightarrow (I, \dot{I}) \in U(1) \times SU(n_H) \subset Sp(2n_H)$ and the symplectic metric $\mathbb{C}_{\alpha\beta}$ reduces to $\mathbb{C}_{IJ} = -\mathbb{C}_{\dot{I}\dot{J}} = \delta_{IJ}$. From equation (30) we see that the constraint (18) for involution is satisfied iff $\mathcal{U}^{1I} \wedge \mathcal{U}^{1\dot{I}} = 0$ that is if, say, the subset $\mathcal{U}^{2I} = (\mathcal{U}^{1I})^*$ of the quaternionic vielbein is set to zero on a submanifold $\mathcal{M}^{KH} \subset \mathcal{M}^Q$.

When this condition is imposed, the submanifold $\mathcal{M}^{KH}$ has dimension at most half the dimension of the quaternionic manifold and the $SU(2)$ connection is reduced to a $U(1)$ connection, whose curvature on $\mathcal{M}^{KH}$ is:

$$\Omega^3|_{\mathcal{M}^{KH}} = i \mathcal{U}^{1I} \wedge \mathcal{U}_{1I} = i \mathcal{U}^{1I} \wedge \overline{\mathcal{U}^{1I}} \quad (31)$$

so that the $SU(2)$ -bundle of the quaternionic manifold is reduced to a $U(1)$ -Hodge bundle for the n_H dimensional complex submanifold spanned by the n_H complex vielbein $\mathcal{U}^{1I}$. The truncation corresponds therefore to select a n_H -complex dimensional submanifold

$\mathcal{M}^{KH} \subset \mathcal{M}^Q$ spanned by the vielbein $\mathcal{U}^{1I}$ and to ask that, on the submanifold, the $2n_H$ extra degrees of freedom are frozen, that is: $\mathcal{U}^{2I}|_{\mathcal{M}^{KH}} = (\mathcal{U}_{1j})^*|_{\mathcal{M}^{KH}} = 0$.

In order to consistently impose the constraint $\mathcal{U}^{2I} = 0$, it is necessary to check its involution:

$$d\mathcal{U}^{2I} = \frac{i}{2}(\omega^1 + i\omega^2) \wedge \mathcal{U}^{1I} + \frac{i}{2}\omega^3 \wedge \mathcal{U}^{2I} - \Delta^I_J \wedge \mathcal{U}^{2J} - \Delta^I_j \wedge \mathcal{U}^{2j} = 0. \quad (32)$$

Eq. (32) implies that we must set to zero, for a consistent reduction, also the “off-diagonal” part of the $Sp(2n_H)$ connection, Δ^I_j . This in turn implies a constraint on the four-fold symmetric tensor $\Omega_{\alpha\beta\gamma\delta}$ appearing in the symplectic part of the quaternionic curvature [2]:

$$\Omega_{IJKL}|_{\mathcal{M}^{KH}} = 0. \quad (33)$$

Finally, let us look at the reduction of the hypermultiplets transformation laws, after imposing (18). They become, after putting $\epsilon_2 = 0$:

$$\mathcal{U}_u^{1I} \delta q^u = \bar{\zeta}^I \epsilon^1; \quad \mathcal{U}_u^{2I} \delta q^u = -\mathbf{C}^{IJ} \bar{\zeta}_j \epsilon_1 \quad (34)$$

$$\delta \zeta_I = i \mathcal{U}_u^{2j} \mathbf{C}_{Ij} \nabla_\mu q^u \gamma^\mu \epsilon^1 + g N_I^1 \epsilon_1 = (\delta \zeta^I)^* \quad (35)$$

$$\delta \zeta_j = i \mathcal{U}_u^{2I} \mathbf{C}_{jI} \nabla_\mu q^u \gamma^\mu \epsilon^1 + g N_j^1 \epsilon_1 = (\delta \zeta^j)^* \quad (36)$$

Equations (34) show that the fermionic partners of the retained scalars $\mathcal{U}^{1I}$ are the spinors ζ_I , while the partners of the scalars $\mathcal{U}^{2I}$ which have to be dropped out are the spinors ζ_j . Consistency of the truncation then imposes that the gauging contribution $g N_j^1$ has to be zero on the reduced $N = 1$ theory.

In the sequel, we just quote the main results on the gauged theory, which are extensively discussed in [2]. The truncation $N = 2 \rightarrow N = 1$ implies, on the gauged theory:

- The D -term of the $N = 1$ -reduced gaugino $\lambda^\Lambda = -2f_i^\Lambda \lambda^{i2}$ is:

$$D^\Lambda = -2f_\alpha^\Lambda W^{\alpha 21} = -2g_{(\Lambda)} (\text{Im} f)^{-1\Lambda\Sigma} \left(P_\Sigma^3(w^s) + P_\Sigma^0(z^i) \right) \quad (37)$$

- The $N = 1$ -reduced superpotential, that is the gravitino mass, is

$$L(z, w) = \frac{i}{2} g_{(X)} L^X \left(P_X^1 - i P_X^2 \right) \quad (38)$$

and is a holomorphic function of its coordinates z^i and w^s .

- The fermion shifts of the $N = 1$ chiral spinors $\chi^i = \lambda^{i1}$ coming from the $N = 2$ gaugini are $N^i \equiv g W^{i11} = 2g^{ij} \nabla_{\bar{j}} \bar{L}$.

- The fermion shifts of the $N = 1$ chiral spinors $\zeta^s = \sqrt{2} P^{I,s} \zeta_I$ coming from $N = 2$ hypermultiplets ($P^{I,s}$ are the scalar vielbein on $\mathcal{M}^{KH}$) are $N^s = -4g_{(X)} k_X^t \bar{L}^X \mathcal{U}_t^{1I} \mathcal{U}_{2j}^s = 2g^{s\bar{s}} \nabla_{\bar{s}} \bar{L}$. Furthermore, the consistency of the truncation of the second gravitino multiplet $\delta\psi_{\mu 2} = 0$ and of the spinors ζ_j in the hypermultiplets sector for the gauged theory gives the gauging constraints:

$$\hat{\omega}_1^2 = 0 \implies g_{(\Lambda)} A^\Lambda \left(P_\Lambda^1 - i P_\Lambda^2 \right) = 0 \implies g_{(\Lambda)} A^\Lambda \left(P_\Lambda^1 - i P_\Lambda^2 \right) = 0 \quad (39)$$

$$S_{12} = 0 \implies g_{(\Lambda)} L^\Lambda P_\Lambda^3 = 0 \implies g_{(X)} L^X P_X^3 = 0 \quad (40)$$

$$\delta \zeta_i = 0 \implies g_{(\Lambda)} k_\Lambda^s \bar{L}^\Lambda = 0 \implies g_{(X)} k_X^s \bar{L}^X = 0 \quad (41)$$

If we call $G^{(2)}$ the gauge group of the $N = 2$ theory and $G^{(1)} \subseteq G^{(2)}$ the gauge group of the corresponding $N = 1$ theory, then we have that the adjoint representation of $G^{(2)}$

decomposes as $Adj(G^{(2)}) \Rightarrow Adj(G^{(1)}) + R(G^{(1)})$, where $R(G^{(1)})$ denotes some representation of $(G^{(1)})$. The gauged vectors of the $N = 1$ theory are restricted to the subset $\{A^\Lambda\}$ generating $Adj(G^{(1)})$ (that is, with the same decomposition of the ungauged theory, we set $\Lambda \rightarrow (\Lambda, X)$, with $\Lambda \in Adj(G^{(2)})$ and $X \in R(G^{(1)})$). The quaternionic Killing vectors of the $N = 2$ theory then decompose as $k_\Lambda^u \Rightarrow \{k_\Lambda^s, k_\Lambda^{\bar{s}}, k_\Lambda^t, k_X^s, k_X^{\bar{s}}, k_X^t\}$. Obviously, we must have that $k_X^s = 0$ since the Killing vectors of the reduced submanifold have to span the adjoint representation of $G^{(1)}$. Viceversa, the Killing vectors with world index in the orthogonal complement, k_Λ^t , must obey $k_\Lambda^t = 0$, while k_X^t are in general different from zero. Indeed, the isometries generated by k_Λ^t would not leave invariant the hypersurface describing the submanifold $\mathcal{M}^{KH} \subset \mathcal{M}^Q$. Since we have found that $k_X^s = 0$, equation (41) is identically satisfied. Eq.s (39) and (40) are satisfied by requiring: $P_\Lambda^1 = P_\Lambda^2 = 0$; $P_X^3 = 0$.

We are left with an $N = 1$ theory coupled to n'_V vector multiplets ($\Lambda = 1, \dots, n'_V$) and $n_C + n_H$ chiral multiplets ($X = 0, 1, \dots, n_C$) with superpotential (38). All the isometries of the scalar manifolds are in principle gauged since the D-term of the reduced $N = 1$ theory depends on $P_\Lambda^0(z, \bar{z}) + P_\Lambda^3(w, \bar{w})$.

References

- [1] J. Strathdee, Int.J.Mod.Phys. **A2** (1987) 273
- [2] L. Andrianopoli, R. D'Auria, S. Ferrara, [hep-th/0110277].
- [3] L.Andrianopoli, M.Bertolini, A.Ceresole, R.D'Auria, S. Ferrara, P.Fré, T.Magri, Nucl.Phys. **B476** (1996) 397; J.Geom.Phys. **23** (1997) 111; R. D'Auria, S. Ferrara, P. Fré, Nucl. Phys. **B359** (1991) 705.
- [4] J. Bagger and E. Witten, Phys.Lett. **B115** (1982) 202; Phys.Lett. **B118** (1982) 103
- [5] E. Cremmer, S. Ferrara, L. Girardello, A. Van Proeyen, Nucl.Phys. **B212** (1983) 231.
- [6] J.A. Bagger, Nucl.Phys. **B211** (1983) 302.
- [7] J.Polchinski, A. Strominger, Phys.Lett. **B388** (1996) 736 [hep-th/9510227]; J. Michel-son, Nucl.Phys. **B495** (1997) 127 [hep-th/9610151]; T.R. Taylor, C. Vafa, Phys.Lett. **B474** (2000) 130 [hep-th/9912152]; P. Mayr, Nucl.Phys. **B593** (2001) 99 [hep-th/0003198]; G.Curio, A. Klemm, D. Luest, S. Theisen, Nucl.Phys. **B609** (2001) 3 [hep-th/0012213]; S.B. Giddings, S. Kachru, J. Polchinski [hep-th/0105097]; K. Behrndt, C. Herrmann, J. Louis, S. Thomas, JHEP **0101** (2001) 011; K. Behrndt, G. Lopes Cardoso, D. Lust, Nucl.Phys. **B607** (2001) 391; G. Dall'Agata [hep-th/0107264]; M.Graña, J.Polchinski, Phys.Rev. **D63** (2001) 026001 [hep-th/0009211]; K. Dasgupta, K. Oh, J. Park, R. Tatar [hep-th/0110050].
- [8] B. de Wit, P.G. Lauwers and A.Van Proeyen, Nucl.Phys. **B255** (1985) 569
- [9] A.Strominger, Comm. Math. Phys. **133** (1990) 163
- [10] L. Castellani, R. D'Auria and S. Ferrara, Phys. Lett. **241B** (1990) 57; Class. Quantum Grav. **7** (1990) 1767.
- [11] L. Andrianopoli, R. D'Auria and S. Ferrara, in preparation.
- [12] A. Ceresole, R. D'Auria and S. Ferrara, Nucl.Phys.Proc.Suppl. **46** (1996) 67-74 [hep-th/9509160].